



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SCIENCE:

PUBLISHED BY N. D. C. HODGES, 874 BROADWAY, NEW YORK.

SUBSCRIPTIONS TO ANY PART OF THE WORLD, \$3.50 A YEAR.

To any contributor, on request in advance, one hundred copies of the issue containing his article will be sent without charge. More copies will be supplied at about cost, also if ordered in advance. Reprints are not supplied, as for obvious reasons we desire to circulate as many copies of *Science* as possible. Authors are, however, at perfect liberty to have their articles reprinted elsewhere. For illustrations, drawings in black and white suitable for photo-engraving should be supplied by the contributor. Rejected manuscripts will be returned to the authors only when the requisite amount of postage accompanies the manuscript. Whatever is intended for insertion must be authenticated by the name and address of the writer; not necessarily for publication, but as a guaranty of good faith. We do not hold ourselves responsible for any view or opinions expressed in the communications of our correspondents.

Attention is called to the "Wants" column. It is invaluable to those who use it in soliciting information or seeking new positions. The name and address of applicants should be given in full, so that answers will go direct to them. The "Exchange" column is likewise open.

THREE SHIPS WITH BERIBERI OUTBREAKS SHOWN TO HAVE HAD EXTENSIVE FORMATION OF CARBONIC OXIDES DURING THE VOYAGE—ANALYSIS OF BERIBERI BLOOD—CONCLUSION THAT BERIBERI IS NOTHING BUT CARBONIC POISONING OF THE BLOOD.

BY ALBERT S. ASHMEAD, M.D., NEW YORK, N.Y.

In *Science*, Nov. 18, 1892, I contributed an account of the outbreak of beriberi on board the bark "H. B. Cann," from Ilo-ilo, Philippine Islands, with a cargo of raw sugar. That cargo fermented during the trip, stifling fumes filled the ship, and the beriberi outbreak was considered the consequence of this state of things.

In an article which will shortly appear in the *Medical News*, entitled "Investigation of the Outbreak of Beriberi on Board the Bark 'Pax' from Ceylon with a Cargo of Graphite," I show that, from the deficient packing of 1,200 tons of graphite, the cargo was exposed to the moist air encountered on a tropical voyage, all but six of a crew of nineteen were stricken with beriberi.

The bark "J. C. Warns," from Java and Macassar, with a cargo of green coffee, arrived in New York June 23, 1890. The captain and three men had died of beriberi. The coffee had been picked and shipped too green. Mr. Tobias, consignee of the cargo, showed me a sample of it; it was charred, carbonized, and almost destroyed. The coffee had fermented. The outbreak of beriberi on a ship from Java, where the coffee has been carbonized, is a regular occurrence. Java coffee owes its value in our market to its color; in order to obtain this color, the captains take their cargoes quite green, which favors a slight fermentation during the trip. Sometimes they go too far; the coffee is too green, and the fermentation too violent; in such cases there is always carbonization; the grains stick together in great masses, and abundant fumes (carbonic gases) fill the ship.

The iron ship "Glenmorag," Captain Currie, 133 days from Colombo, Ceylon, with 1,100 tons of graphite on board, 800 tons of cocoa-nut oil, etc., arrived July 17 in New York. This ship, loaded in Colombo alongside the "Pax" (mentioned above), travelled the same course, at an interval of two weeks. She lies now at the Atlantic dock, in Brooklyn, again alongside the "Pax." She had no beriberi outbreak. From her first mate I have the following information:—

Crew, 28 men; captain's wife and two children on board; or all, 31.

She is a Scotch Glasgow boat, and the crew is English and Scotch. Before taking this cargo, this ship had carried from Barry Dock, near Cardiff, a cargo of coal to Buenos Ayres, South America, and taken a ballast of sand to Colombo, Ceylon. Before these trips she had been in the wheat trade from Tacoma, Washington, to Havre, France. She is remarkably dry, and the cleanest ship one would wish to see. I went down her hold and examined every part of it; there is not a smut nor a stain anywhere about it. The iron part is especially clean: no trace of incrustation of carburetted iron, which might have indicated the action of hot moist

air on the carbon. None of the barrels containing the graphite was broken. The packing was exceedingly good, the dunnage consisted of sticks and cocoa-nut hulls, so that it was impossible for the barrels to roll and break, and thus expose their contents to the action of the air.

The diet bill was about the same, or even poorer, than that of the "Pax." Nine casks of salted beef and seven barrels of pork were consumed during the trip. Fresh beef (tinned) three times a week, one-half pound to a man, and a half pound of salt meat on the same days; other days a full pound of salt meat a day. One-half pound of rice for each man on Saturdays; no vegetables except onions with the soup three times a week. The ship being Scotch, oatmeal made part of its fare for two and a half months after starting, when it ran out. No sickness whatever during the voyage. One death by accident. The captain attributes the good condition of his cargo and his crew to the change of winds and cooler weather which he enjoyed from the Cape of Good Hope to the North Atlantic. His log is indeed very different from that of the "Pax."

In *Science*, vol. xxii., No. 545, p. 16, Venable states that "the metallic carbides are usually formed by the action of intense heat upon the metal in the presence of carbon. The form of this carbon is capable of being greatly varied. Graphite, amorphous carbon, and many hydrocarbons, may be used. The heat of the ordinary furnace is sufficient to form the carbides of the metals already mentioned, zinc, copper, and notably iron. All of these carbides, under certain conditions, give off their carbon in the form of hydrocarbons. The same smell can be detected in all during their decomposition. In some cases, as iron and zinc, the decomposition is caused by the action of an acid. The carbides of the earths (*of which graphite is one, in conjunction with iron*) decompose in moist air, and more rapidly in water."

I may point again here to those broken barrels of the "Pax," which exposed the carbon to the influence of tropical air.

I have examined microscopically the blood of four of the sufferers of the "Pax," and obtained the following results: Captain Geeseicke, sick since May 16 with beriberi; 800 diameters, $\frac{1}{12}$ inch oil-immersion objective; red discs, irregular in outline, congregated in masses, with ragged edges, not inclined to form rouleaux; quite plastic; colored streaks or rays of pink and red, showing the presence of biliary matter, biliverdin crystals; black spores, not free but entangled in the hummocks of corpuscles. It may be noted that the oedema of this patient's legs only left him two days before this examination.

Henry Oelrichs (second mate), German, 27 years old. Has been fourteen days sick with beriberi. Examination of the blood: 500 diameters, $\frac{1}{8}$ inch objective; red corpuscles, very plastic, aggregated in hummocks. Many black spores are seen floating about, free in motion. Fibrine in excess, light in texture, and lumpy. Blood very thick, syrupy, and plastic. No motion, showing want of circulation. Excess in the coloring matter. This same case examined: 900 diameters, $\frac{1}{16}$ inch objective immersion lens, shows excess of fibrine in ropes, biliary matter in great excess; no crystalline formations; blood quickly oxidizes and forms a solid mass. The black spores above mentioned are quickly held by the fibrine; the red discs are distended, bladder-shaped, and have very ragged edges. The meniscus-shape is lacking, there being great irregularity in outline and color, some are even square-shaped. Some discs have an excess of color, some are very pale. On the edges of the corpuscular mass the color quickly disappears, in consequence of rapid coagulation.

Isaac Heggund, a Swede, 27 years old, has had beriberi since crossing the equator, six weeks ago. Legs are now very thin, but still some soreness remains; knee reflexes still lost. First sound of heart prolonged. Microscopical examination of the blood, 900 diameters, $\frac{1}{16}$ inch objective, shows rouleaux well formed, no spores, no filaments, slight feverish condition shown by spiculated outlines of some of the red corpuscles. Fibrine is assuming a normal form, showing meshes very regular; no distension of red corpuscles.

Emil Jensen, a German, 19 years old, sixteen days sick with beriberi. Black spores in active motion and very plentiful; freely scattered in the field of observation; circulation very torpid; fibrine